

DIRECTIONS FOR USE

WEED TABLE

NOTE: Where weeds are to be sprayed in a CROP or PASTURE, use only the rates given for the crop in the table below. In most cases this will give control, however some hard to kill weeds eg. Rumex spp. (docks) and Polygonum spp. (wireweed, climbing buckwheat) or those in advanced stages of growth may only be suppressed. The product should only be used in those States where a rate or range of rates is indicated for the particular weeds listed.

APPLICATION RATE PER HECTARE

Weeds Controlled	Crop						Pasture – Non Legumes		CRITICAL COMMENTS
	VIC	SA	TAS	NSW	QLD	WA	VIC	NSW, SA, QLD, WA ONLY	
Amaranthus spp	-	-	-	900mL	-	-	-	-	
Angled Onion	-	-	-	-	-	-	3.7L	900mL-1.9L	Spray when buds forming or early flowering
Apple of Sodom	-	-	-	-	-	-	-	3.3-3.7L	
Bathurst Burr	-	-	-	900mL	-	-	1.9-3.7L	1.9-3.7L	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Black Knapweed	-	-	-	-	-	-	3.7L	-	Spray before flowering. DO NOT cultivate these infestations.
Buffalo Burr	-	-	-	-	-	-	-	900mL-1.3L (not Qld & WA)	Spray before seedling to pre-flowering. Use higher rate as plant matures
California Burr	-	-	-	900mL	-	-	1.9-3.7L	X	Spray from seedling to pre-flowering. Use higher rate as plant matures
Caltrop	-	-	-	700 - 900mL	-	-	1.9-3.7L	-	Spray from seedling to pre flowering. Use higher rate as plants mature
Cape Tulip	-	-	-	-	-	1.3L	3.7L	1.9-3.7L	Spray before flowering
Capeweed	900mL	900mL	900mL	600 - 900mL	-	-	-	2.8-3.7L	Spray up to rosette stage
Charlock	460mL	460mL	900mL	460mL	-	-	-	900mL	Spray up to rosette stage
Clover	-	-	-	700 - 900mL	-	-	-	900mL	
Colocynth	-	-	-	-	-	-	3.7L	-	Spray at seedling stage only
Deadnettle	-	-	-	900mL	-	-	-	-	
Devil's Claw	-	-	-	900mL	-	-	1.5L	X	Spray prior to pods forming

Weeds Controlled	Crop						Pasture – Non Legumes		
	VIC	SA	TAS	NSW	QLD	WA	VIC	NSW, SA, QLD, WA ONLY	CRITICAL COMMENTS
Dock	900mL	900mL	-	-	900mL	900mL	-	1.9-2.8L	Spray at rosette stage to kill top growth only
Fat Hen	-	-	-	460-900mL	-	-	-	-	
Flatweed	-	-	-	900mL	-	-	-	2.8-3.7L	
Fumitory - red	-	900mL	-	900mL	-	-	-	2.8-3.7L	Spray up to rosette stage
Fumitory – white	900mL	460mL	-	900mL	-	-	-	2.8-3.7L	Spray up to rosette stage
Galvanised Burr	-	-	-	-	-	-	5.3L	5.3L (not Qld & WA)	Spray from seedling to pre-flowering
Goosefoots	-	-	-	900mL	-	-	-	-	
Hard Head or Russian Knapweed	-	-	-	-	-	-	3.7-5.9L	-	Spray before flowering
Hogweed/ Wireweed	900mL	900mL	-	900mL	900mL	-	-	1.3-1.9L (Not SA)	Spray up to rosette stage
Hoary Cress/ Whiteweed	-	900mL	900mL	900mL	-	-	1.9-3.7L	1.9-2.4L	Spray from late rosette to pre- flowering
Horehound (seedlings)	-	900mL	-	-	-	950mL	-	1.9-3.7L	Late Autumn to early Spring
Iron Weed, Corn Cromwell	-	-	-	900mL	-	-	-	1.3-1.9L	
Khaki Weed	-	-	-	-	-	-	-	900mL-1.3L (not SA)	Spray young seedlings only
Lincoln Weed	-	900mL	-	-	-	-	-	-	Autumn spray before sowing improves control
London Rocket	-	-	-	-	-	650mL	-	1.8-2.8L (WA only)	
Lupins	900mL	-	-	460 - 900mL	-	-	-	-	Spray up to rosette stage
Melilotus/Hexham Scent	900mL	900mL	-	-	900mL	-	-	1.3-1.9L	Spray up to rosette stage
Melons - camel, paddy	-	-	-	460 - 900mL	-	-	-	-	
Mustards	370mL	260 - 900mL	900mL	460 - 900mL	700mL	700mL	3.7L	1.9-2.8L	Spray up to rosette stage

Weeds Controlled	Crop						Pasture – Non Legumes		
	VIC	SA	TAS	NSW	QLD	WA	VIC	NSW, SA, QLD, WA ONLY	CRITICAL COMMENTS
Mexican Poppy	-	-	-	900mL	-	950mL	-	900mL-1.3L (1.3 – 1.7 L WA only)	Spray rosette stage and before flowering.
Mintweed	-	-	-	900mL	700mL	-	-	900mL-1.3L	Spray active seedlings only.
Muskweed	900mL	-	-	-	-	-	-	-	Spray up to early rosette stage
New Zealand Spinach	-	-	-	900mL	-	-	-	-	
Noogoora Burr	-	-	-	900mL	-	-	1.9-3.7L	1.9-3.7L	Spray seedling to pre-flowering.
Nut Grass	-	-	-	-	-	-	3.7-5.9L	-	Spray within 4 weeks of foliage emergence, repeat spray necessary
Paterson's Curse	-	-	-	900mL	-	950mL	1.9-3.7L	900mL-1.9L (1.3-1.7L WA only)	Spray seedling to rosette stage
Poppy, Wild	460mL	-	-	-	-	-	-	2.4-3.3L	Spray up to rosette stage
Ragwort	-	-	-	-	-	-	3.7L	3.7L	Spray at rosette to cabbage stage
Rapeseed	900mL	-	-	460 - 900mL	-	-	-	-	Spray up to rosette stage
Rapistrum	-	-	-	-	-	650mL	-	950mL (WA only)	
Rough Poppy	-	460mL	-	460 - 900mL	-	-	-	900mL	Spray young seedlings only
St Johns Wort	-	-	-	-	-	-	3.7-5.9L	3.7-5.3L	Spray before flowering. Spray before plants are 40cm high
Safflower	-	-	-	460 - 900mL	-	-	-	-	
Sand Mustard/ Sand Rocket	-	-	-	-	-	-	3.7L	-	Spray before flowering
Shepherd's Purse	-	-	-	900mL	-	-	-	-	
Silverleaf	-	-	-	-	-	-	3.7L	-	Spray at flowering. Fallow land: controls top growth only
Nightshade	-	-	-	-	-	-	-	-	
Skeleton Weed	-	900mL	-	900mL	-	-	3.7L	1.3-1.9L	Spray rosettes before aerial growth commences

Weeds Controlled	Crop						Pasture – Non Legumes		
	VIC	SA	TAS	NSW	QLD	WA	VIC	NSW, SA, QLD, WA ONLY	CRITICAL COMMENTS
Stingless Nettle (Deadnettle)	-	900mL	90mL	-	-	-	-	2.4-2.8L	
Stinging Nettle	900mL	-	-	-	-	-	-	-	Spray up to rosette stage
Stinkwort	-	-	-	900mL	-	-	1.9-3.7L	1.9-3.7L	Spray younger plants. Use higher rate as plants mature.
Storksbill/Erodium	-	-	-	900mL	-	-	-	-	
Sunflower seedlings	900mL	-	-	460-900mL	700mL	-	-	-	Spray multiple leaves
Thistles									
- Golden	-	-	-	-	-	-	3.7L	3.7L	Spray at rosette stage
- Nodding	-	-	-	-	-	-	3.7L	1.3-1.9L	Spray rosette to pre-flowering stage
- Saffron	700mL	900mL	-	460-900mL	900mL	900mL	900mL-1.9L	900mL-2.8L	Spray up to rosette stage
- Sheep	-	-	-	-	-	950mL	-	950mL-3.7L (WA only)	
- Slender, Shore	-	-	900mL	900mL	-	X	1.9-3.7L	900mL-3.7L	Spray at rosette stage
- Soldier	-	-	-	-	-	-	3.7L	-	Spray at rosette stage
- Spear	900mL	-	900mL	-	-	-	900mL-2.8L	1.3-2.4L	Spray at seedling to rosette stage. Use higher rate as plants mature (pastures)
- Stemless	-	-	-	-	-	-	3.7L	2.8-3.7L	Spray rosette stage to flowering
- St Barnaby's	-	-	-	-	-	-	-	1.3-1.9L	
- Star	-	-	-	900mL	-	-	1.9-3.7L	X	Spray seedling to rosette stage. Use higher rate as plants mature.
- Variegated	-	-	900mL	460-900mL	700mL	-	900mL-2.8L	900mL-3.7L	Spray at rosette stage. Can cause stock poisoning
Thornapple	-	X	-	460-900mL	-	-	3.7L	900mL-1.9L	Spray at seedling stage.
Tree Hogweed	900mL	-	-	-	-	-	-	-	Spray up to rosette stage
Turnip Weed	-	460mL	-	460-900mL	460mL	700mL	-	900mL	Spray seedlings only
Vetches Tares	900mL	700mL	900mL	-	-	-	-	-	

Weeds Controlled	Crop						Pasture – Non Legumes		
	VIC	SA	TAS	NSW	QLD	WA	VIC	NSW, SA, QLD, WA ONLY	CRITICAL COMMENTS
Wards Weed	-	460mL	-	-	-	-	-	-	Spray at seedling stage.
Wild Cabbage	900mL	-	-	-	-	-	-	-	Spray up to rosette stage
Wild Garlic	-	-	-	-	-	-	7.5L	-	Suppresses aerial growth only.
Wild Mignonette	-	-	-	-	-	950mL	3.7L	-	Spray up to rosette stage
Wild Mustard	-	-	-	-	-	650mL	-	1.8-2.8L (WA only)	
Wild Radish	900mL	900mL	900mL	460-900mL	900mL	650mL	-	900mL (950mL WA only)	Spray up to rosette stage.
Wild Sage	-	-	-	-	-	-	-	2.8-3.7L	
Wild Teasel	-	-	-	-	-	-	1.9-3.7L	-	Spray at rosette stage. Use the higher rate as plants mature
Wild Turnip	240mL	260mL	900mL	460-900mL	-	450mL	-	900mL (950mL WA only)	Spray up to rosette stage